

ON GENERALIZED STONE'S THEOREM

MASSOUD AMINI

ABSTRACT. It is known that the generator of a strictly continuous one parameter unitary group in the multiplier algebra of a C^* -algebra is affiliated to that C^* -algebra. We show that under natural non degeneracy conditions, this self adjoint unbounded operator lies indeed in the (unbounded) multiplier algebra of the Pedersen's ideal of the C^* -algebra.

1. INTRODUCTION

One of the main objectives of the Heisenberg formulation of the Quantum Mechanics is to give appropriate models for the *commutation relations*, the most famous one of which being

$$[P, Q] = i\hbar I$$

where $\hbar$ is the *Planck constant* and P, Q are the *quantum position* and *quantum momentum*. It was known from the beginning that bounded linear operators can not satisfy such a relation (convince yourself by checking this for matrices where you have a *trace* for free!). In particular this does not happen in a C^* -algebra. Although projective limit of C^* -algebras can include unbounded operators, this can *not* happen in a projective limit also. (Just recall that some of their quotients are C^* -algebras [Ph88b]). One classical trick is to replace this type of relation by a stronger commutation property, which in this case is

$$U_t Q U_{-t} = e^{i\hbar t} Q$$

where $U_t = e^{itH}$, for some closed operator H . It has been shown in [HQV] that any strictly continuous one parameter group of unitaries (U_t) in the multiplier algebra of a C^* -algebra A_0 has a generator H (a version of Stone's theorem) which could be chosen to be in A_0^η (the set of densely defined linear operators on A_0 which are affiliated with A_0 [Wr91]). Here we show that indeed H could be chosen more specifically. Let $A_{00} = K(A_0)$ be the Pedersen's ideal of A_0 and $\Gamma(A_{00})$ be the topological algebra of (unbounded) multiplier on A_{00} [LG]. We show that H could be chosen to be in $A = \Gamma(A_{00})$.

2. STONE'S THEOREM

We start with some lemmas from [Wr91] in which we replace A_0^η with $A = \Gamma(A_{00})$. We keep the notations of the above paragraph all over the paper. We also freely use the notations and terminology of [Wr91]. All morphisms are supposed to be non degenerate, and following [Wr 91] we use Mor to denote the set of morphisms.

1991 *Mathematics Subject Classification*. Primary 46L05; Secondary 47D03.

Key words and phrases. Stone's theorem, Pedersen's ideal, unbounded multipliers. Partially supported by an IPM grant.

The following lemma has been already proved in [Am], but we bring the proof here for the sake of completeness.

Lemma 2.1. *Let A_0 and B_0 be C^* -algebras. Let $\phi_0 \in \text{Mor}(A_0, B_0)$ be strictly non degenerate (that is $B_{00} \subseteq \phi_0(A_{00})B_0$). Then ϕ_0 extends uniquely to a morphism $\phi \in \text{Mor}(A, B)$ such that $\phi(A_{00})B_0$ is a core for $\phi(T)$ and $\phi(T)(\phi_0(a)b) = \phi_0(Ta)b$ for all $a \in A_{00}$, $b \in B_0$, and $T \in A = \Gamma(A_{00})$.*

Proof Since ϕ_0 preserves the spectral theory, $\phi_0(A_{00}) \subseteq B_{00}$. On the other hand, $\phi_0(A_{00})$ is clearly a dense ideal of $\phi_0(A_0)$. Therefore $B_0\phi_0(A_{00})B_0$ is an ideal of B_0 which is dense in $B_0\phi_0(A_0)B_0$. Now if ϕ_0 is non degenerate then $\phi_0(A_0)B_0$ is dense in B_0 , so $B_0\phi_0(A_0)B_0$ is dense in $B_0B_0 = B_0$, i.e. $B_0\phi_0(A_{00})B_0$ is a dense ideal of B_0 and so contains B_{00} . But $B_0\phi_0(A_{00})B_0 \subseteq B_0B_{00}B_0 \subseteq B_{00}$, hence the equality holds. If ϕ_0 is strictly non degenerate then $\phi_0(A_{00})B_0 \supseteq B_{00}$. The converse inclusion follows from the fact that $\phi_0(A_{00}) \subseteq B_{00}$. Hence $\phi_0(A_{00})B_0 = B_{00}$. Now the right hand side is self adjoint and the adjoint of the left hand side is $B_0\phi_0(A_{00})$, hence $B_0\phi_0(A_{00}) = B_{00}$. $\square$

Now each $T \in A = \Gamma(A_{00})$ could be considered as an element of A_0^η and ϕ_0 also extends to $\tilde{\phi} : A_0^\eta \rightarrow B_0^\eta$ by [Wr91, 1.2]. But there is no ambiguity as we have

Lemma 2.2. *With the above notations, $\tilde{\phi}(T) = \phi(T)$. In particular we have $\phi(z_T) = z_{\phi(T)}$.*

Proof The first statement follows from above lemma and the facts that by assumption $\phi_0(A_{00})B_0 \supseteq B_{00}$ and $\tilde{\phi}(T)(\phi_0(a)b) = \phi_0(Ta)b$ ($a \in D(T) \supseteq A_{00}$, $b \in B_0$) [Wr91, 1.2]. The second statement is proved for $\tilde{\phi}$ in [Wr91, 1.2], and follows from the first for ϕ . $\square$

Proposition 2.1. *Let A_0 be a C^* -algebra and $T \in A$ be self adjoint. Let $z_T \in M(A_0)$ be the z -transform of T . Then $\sigma(z) \subseteq [-1, 1]$. Assume that*

$$\text{span}\{f(z_T)a : f \in C_0(-1, 1), a \in A_0\} \supseteq A_{00}.$$

Then there is a unique $\phi = \phi_T \in \text{Mor}(C(\mathbb{R}), A)$ such that $\phi_T(id) = T$. Moreover $\phi(C_0(\mathbb{R}))A_0 \supseteq A_{00}$ and $\phi_T(z) = z_T$, for $z(t) = t(1 + t^2)^{\frac{1}{2}}$, $t \in \mathbb{R}$.

Proof $z_T \in M(A_0)$ is self adjoint and $\|z_T\| \leq 1$, so its spectrum $\sigma(z_T)$ is contained in $[-1, 1]$. The same is true for $z \in C_b(\mathbb{R})$. We use the continuous functional calculus to show the uniqueness. If $\phi(id) = T$ then $\phi(z) = z_T$ and $\phi(f \circ z) = f(z_T)$, for each $f \in C_b(-1, 1) = C[-1, 1]$. But each element of $C_0(\mathbb{R})$ is of the form $f \circ z$, where $f \in C_0(-1, 1)$ and the uniqueness follows.

For the existence, let's define $\phi_0 : C_0(\mathbb{R}) \rightarrow M(A_0)$ by $\phi_0(f \circ z) = f(z_T)$. Then ϕ_0 is clearly a $*$ -homomorphism. Also by assumption, $\phi_0(C_0(\mathbb{R})A_0) \supseteq A_{00}$ which means that ϕ_0 is strictly non degenerate. By [Am, Theorem 3.4], ϕ_0 extends (uniquely) to some $\phi \in \text{Mor}(C(\mathbb{R}), \Gamma(A_{00}))$. Now $\phi(f \circ z) = f(z_T)$, for each $f \in C_0(-1, 1)$. In particular, for $f(t) = t(1 - t^2)^{\frac{-1}{2}}$, we get $f \circ z = id$ and so $\phi(id) = f(z_T) = T$. $\square$

Let ϕ_0 and ϕ be as in the proof of the above proposition. Then ϕ is not injective in general. But we can do the classical trick to make it injective. Consider $\ker(\phi_0) = \{f \in C_0(\mathbb{R}) : f = 0 \text{ on } \sigma(T)\}$. Then $C_0(\mathbb{R})/\ker(\phi_0) = C_0(\sigma(T))$ and the corresponding quotient map identifies with the restriction map $\pi : C_0(\mathbb{R}) \rightarrow C_0(\sigma(T))$. Let $id_{\sigma(T)} = \pi(id)$, then we have

Proposition 2.2. *Let A_0 be a C^* -algebra and $T \in \Gamma(A_{00})$, then there is a unique embedding $\psi = \psi_T \in \text{Mor}(C(\sigma(T)), \Gamma(A_{00}))$ such that $\psi_T(id_{\sigma(T)}) = T$.*

Proof Since π is onto, there is a function ψ_0 such that $\psi_0\pi = \phi_0$. Then ψ_0 is a $*$ -homomorphism. To see that it is strictly non degenerate it is enough to observe that $\psi_0(C_0(\sigma(T)))A_0 = \phi_0(C_0(\mathbb{R}))A_0 \supseteq A_{00}$. Therefore it extends to a morphism $\psi \in \text{Mor}(C(\sigma(T)), \Gamma(A_{00}))$.

Next let us show that ψ is one-one. It is clear that ψ_0 is one-one (since $\psi_0 = \phi_0\pi$ where $\pi(\ker \phi_0) = \{0\}$). Take any $F \in C(\mathbb{R})$ such that $\psi(F) = 0$, then $0 = \psi(F)\psi_0(f)a = \psi_0(Ff)a$, for each $f \in C_0(\sigma(T))$ and $a \in A_0$. This means that $\psi_0(Ff) \in M(A_0)$ multiplies A_0 into 0, i.e. $\psi_0(Ff) = 0$. Hence $Ff = 0$ for each $f \in C_0(\sigma(T))$, and so $F = 0$. $\square$

Now we are prepared to prove the generalization of Stone's theorem. Let A_0 be a C^* -algebra, $A_{00} = K(A_0)$ its Pedersen's ideal, and $A = \Gamma(A_{00})$ be the algebra of (unbounded) multipliers of A_{00} [LT]. For each $t \in \mathbb{R}$ consider the function $e_t \in C(\mathbb{R})$ defined by $e_t(s) = \exp(its)$ ($s \in \mathbb{R}$). Let $h \in A = \Gamma(A_{00})$ and $U_t = \phi_h(e_t) = \exp(ith)$. Then $(U_t)_{t \in \mathbb{R}}$ is a one parameter strictly continuous unitary group in $M(A_0)$ (the strict continuity follows from the fact that ϕ_h is non degenerate). Moreover, if $h \in b(\Gamma(A_{00})) = M(A_0)$, then this is also norm continuous. Conversely each strictly continuous unitary group in $M(A_0)$ is of this form, for some $h \eta A_0$ [HQV]. Here we want h to be actually in $\Gamma(A_{00})$. Clearly for this to happen, we would need to put some condition on the unitary group. This is the content of the following result. The proof is quite similar to [HQV, 2.1]. Here we only sketch those parts of the proof which have to be modified. But first a definition.

Definition 2.1. *Let A_0 be a C^* -algebra and $(U_t)_{t \in \mathbb{R}}$ be a one parameter strictly continuous unitary group in $M(A_0)$. Let's define $\alpha : L^1(\mathbb{R}) \rightarrow M(A_0)$ by $\alpha(f) = \int_{\mathbb{R}} f(t)U_t dt$, where*

$$\left(\int_{\mathbb{R}} f(t)U_t dt \right) x = \int_{\mathbb{R}} f(t)U_t x dt \quad (x \in A_0)$$

is in the Bochner sense. This extends to a $$ -homomorphism $\alpha \in \text{Mor}(C^*(\mathbb{R}), A_0)$ [HQV]. We say that (U_t) is (strictly) non degenerate, if the morphism α is (strictly) non degenerate (cf. [Am]).*

Theorem 2.1. (Generalized Stone's Theorem) *Let A_0 be a C^* -algebra and $(U_t)_{t \in \mathbb{R}}$ be a one parameter strictly continuous strictly non degenerate unitary group in $M(A_0)$. Then there exists a self adjoint $h \in \Gamma(K(A_0))$ such that $U_t = \exp(ith)$ for $t \in \mathbb{R}$. Moreover, if $(U_t)_{t \in \mathbb{R}}$ is norm continuous, then $h \in M(A_0)$.*

Proof Define $\alpha \in \text{Mor}(C^*(\mathbb{R}), A_0)$ as above. Then α is strictly non degenerate, and so it extends to a morphism of the corresponding pro- C^* -algebras, which we still denote it by $\alpha \in \text{Mor}(\Gamma(K(C^*(\mathbb{R}))), \Gamma(K(A_0)))$ (see the discussion before Theorem 3.2 in [Am]). Also it is well known that the Fourier transform $\mathcal{F} : L^1(\mathbb{R}) \rightarrow C_0(\mathbb{R})$ extends to an isomorphism $\mathcal{F} \in \text{Mor}(C^*(\mathbb{R}), C_0(\mathbb{R}))$ with $\mathcal{F}(e_t) = \lambda_t$, where λ is the left regular representation of $\mathbb{R}$. Since $\mathcal{F}$ is surjective we can extend it to an isomorphism $\mathcal{F} \in \text{Mor}(\Gamma(K(C^*(\mathbb{R}))), C(\mathbb{R}))$ [LT]. Also it is clear that $\alpha(\lambda_t) = U_t$ ($t \in \mathbb{R}$). Define $h = (\alpha \circ \mathcal{F}^{-1})(id) \in \Gamma(A_{00})$. This is self adjoint and by the uniqueness part of Proposition 1.2 we have $\phi_h = \alpha \circ \mathcal{F}^{-1}$. Now $U_t = \alpha(\lambda_t) =$

$\alpha(\mathcal{F}^{-1}(e_t)) = \phi_h(e_t) = \exp(ith)$, which finishes the proof of the first part of the theorem.

Now, for each $t \in \mathbb{R}$ we get

$$\begin{aligned} \|U_t - 1\| &= \|\phi_h(e_t - 1)\| = \|\psi_h \pi(e_t - 1)\| = \|\pi(e_t - 1)\| \\ &= \sup\{|e^{it\lambda} - 1| : \lambda \in \sigma(h)\}. \end{aligned}$$

So, if the unitary group is norm continuous, then $\sigma(h)$ is bounded. Hence $h \in b(\Gamma(K(A_0)))$ [Ph88b]. But $b(\Gamma(K(A_0))) = M(A_0)$ [Ph88a], so $h \in M(A_0)$ and the proof is complete. $\square$

Next, following [HQV], we show that h can be found by *differentiating* the unitary group $(U_t)_{t \in \mathbb{R}}$.

Proposition 2.3. *Let A_0 be a C^* -algebra, $(U_t)_{t \in \mathbb{R}}$ a one parameter strictly continuous unitary group in $M(A_0)$, and h a self adjoint element of $\Gamma(K(A_0))$ such that $U_t = \exp(ith)$ for $t \in \mathbb{R}$. Define the (unbounded) operator $H : D(H) \subseteq A_0 \rightarrow A_0$ by*

$$\begin{aligned} (1) \quad D(H) &= \{a \in A_0 : t \mapsto U_t a \text{ is } C^1\} \\ (2) \quad Ha &= \frac{d}{dt}|_{t=0} U_t a = \lim_{t \rightarrow 0} (U_t a - a)/t \quad (a \in D(H)). \end{aligned}$$

Then $h \subseteq iH = i \frac{d}{dt}|_{t=0} U_t$.

Proof Using minimality of A_{00} we have $A_{00} \subseteq \sqrt{1 - z_h} A_0$. Now this last set is contained in $D(H) = D(iH)$ [HQV, 2.2]. The fact that $h(x) = iH(x)$, for each $x \in A_{00}$, follows from the calculations of [HQV, 2.2]. $\square$

This in particular shows the uniqueness of the element $h \in \Gamma(A_{00})$ for which $U_t = \exp(ith)$ ($t \in \mathbb{R}$). We call this element the *infinitesimal generator* of the one parameter group $(U_t)_{t \in \mathbb{R}}$.

3. ELEMENTS AFFILIATED WITH GROUP C^* -ALGEBRA

The group C^* -algebras are important objects in the theory of quantum groups. One reason is that they are *dual objects* to continuous functions. When the underlying group is not discrete (non compact case), one would expect some unbounded elements to come into the play. These can not belong to the group C^* -algebra, but they are usually affiliated with it. The problem of finding all elements affiliated with $C^*(G)$ is open in general. Some attempts are done to find them in the case that G is a Lie group [Wr91]. In this case we know that the elements of the corresponding Lie algebra, considered as differential operators, are affiliated with the group C^* -algebra. The question of whether all elements of the universal enveloping algebra of G are affiliated with $C^*(G)$ was left open. In this section we give a partial answer to this question using pro- C^* -algebras.

Let G be a Lie group, and $\pi : G \rightarrow B(\mathcal{X})$ be a (strongly continuous) representation of G on a Banach space $\mathcal{X}$. We say that $x \in \mathcal{X}$ is a C^∞ -vector (analytic vector, respectively) of π if the map $g \mapsto \pi(g)x$ from G to $\mathcal{X}$ is a C^∞ (analytic, respectively) function. We denote the set of all such elements $x \in \mathcal{X}$ by $D^\infty = D^\infty(\pi)$ ($D^\omega(\pi)$, respectively). Then this is a dense subset of $\mathcal{X}$. Indeed L. Gårding showed that if $\phi \in C_{00}^\infty(G)$ and $\pi(\phi)x = \int_G \pi(t)x\phi(t)dt$ ($x \in \mathcal{X}$), then $D_{00}^\infty = \pi(C_{00}^\infty(G))\mathcal{X} \subseteq \mathcal{X}$ is dense. This is called the *Gårding domain* of π . Let $\mathcal{G} = L(G)$ be the Lie algebra of

G , i.e. the set of all left invariant vector fields on G (at identity). For each $X \in \mathcal{G}$, define

$$\pi(X)x = \lim_{h \rightarrow 0} \frac{\pi(\exp hX)x - x}{h} \quad (x \in D^\infty)$$

Then $\pi(X) : D^\infty \subseteq \mathcal{X} \rightarrow D^\infty \subseteq \mathcal{X}$ is a densely defined unbounded operator on $\mathcal{X}$, and it is *skew symmetric* if π is unitary (and $\mathcal{X}$ a Hilbert space).

Harish-Chandra noticed that D^∞ could have a subspace D such that $\pi(X)D \subseteq D$ but $\pi(g)\bar{D} \not\subseteq \bar{D}$. Therefore he suggested replacing D^∞ with D^ω . He showed that D^ω is dense in $\mathcal{X}$ for certain representations of a semisimple Lie group [Har]. P. Cartier and J. Dixmier gave a proof for all unitary representations [CD2], and E. Nelson used a generalization of the fundamental solution of the heat equation on Lie groups to prove this for an arbitrary representation. He also used analytic vectors to give a sufficient condition for a representation of a Lie algebra to be induced by a unitary representation of the Lie group [Nel]. Later I. Segal showed that this is equivalent to the *complete positivity* of the representation with respect to an appropriate cone [Seg].

We are mainly interested in the case of the universal representation. For any locally compact group G , the universal representation $u : G \rightarrow M(C^*(G))$ is determined by the following universal property: For each C^* -algebra A and each representation $\pi : G \rightarrow A$, there is a unique $\tau \in Mor(C^*(G), A)$ such that $\pi = \tau u$. Then u is continuous and open with respect to the strict topology. Now let G be a Lie group and let $\mathcal{G}$ be its Lie algebra of dimension N with a basis $X_1, X_2, \dots, X_N$. Then the *universal enveloping algebra* $\mathcal{U} = \mathcal{U}(\mathcal{G})$ of $\mathcal{G}$ is an $*$ -algebra under $X^* = -X$, $X \in \mathcal{G}$. Elements of $\mathcal{U}$ are *differential operators* on G commuting with the right translations: Take $D^\infty = D^\infty(u)$, then each $X \in \mathcal{U}$ defines

$$\begin{aligned} du(X) : D^\infty \subseteq C^*(G) &\rightarrow B(H_u) \\ du(X)a &= Xu(g)a|_{g=e} = \lim_{t \rightarrow 0} \frac{1}{t}(u(\exp tX)a - a) \end{aligned}$$

which is a closable operator whose closure is simply denoted by $X : D^\infty \subseteq C^*(G) \rightarrow C^*(G)$. For the reasons explained in the beginning of this section, we are interested in elements affiliated with $C^*(G)$. We know that for $X \in \mathcal{U}$, if the differential equation $X^*Xf = -f$ has only trivial bounded C^∞ -solution, then $X, X^* \eta C^*(G)$ [Wr95,2.1]. It is known that all the elements of $\mathcal{G}$ and also the *elliptic operator* $\Delta = -\sum_1^N X_i^2$, where X_i 's form a basis of $\mathcal{G}$, are affiliated with $C^*(G)$ [Wr95]. The X_i 's are skew self-adjoint, where as Δ is self-adjoint and positive (yes positive!).

To look more carefully at the above problem, one has to distinguish between the right and left multipliers. To each $g \in G$ we associate a bounded operator $u(g) \in B(H_u)$. We can view this as a bounded multiplier on $C^*(G)$, by considering it as an element of $C^*(G)^{**}$. This means that we put $u(g) = (S_g, T_g)$, where S_g and T_g are bounded extensions of the mappings

$$S_g(u(f)) = u(\delta_g * f) \quad \text{and} \quad T_g(u(f)) = u(f * \delta_g) \quad (f \in L^1(G))$$

Now consider the following

$$D_\ell^\infty = \{a \in C^*(G) : g \mapsto S_g(a) \text{ is a } C^\infty \text{ map}\}$$

and define

$$d_\ell u(X) : D_\ell^\infty \subseteq C^*(G) \rightarrow C^*(G)$$

$$d_\ell u(X)a = \lim_{t \rightarrow 0} \frac{1}{t} (S_{\exp(tX)}(a) - a) \quad (a \in D_\ell^\infty)$$

then D_ℓ^∞ is a dense right ideal of $C^*(G)$. In order to connect this with the above problem, we need to define the one sided version of some of the previous notions.

Definition 3.1. Let J be a right ideal in a C^* -algebra A_0 . Then a left multiplier of J is a linear map $S : J \rightarrow J$ such that $S(ax) = S(a)x$ ($a \in J, x \in A_0$). The set of all such maps is denoted by $\Gamma_\ell(J)$. A right multiplier is defined similarly and we denote the set of all right multipliers by $\Gamma_r(J)$.

There is a natural left version of the κ -topology for $\Gamma_\ell(J)$: A net $(S_\alpha) \in \Gamma_\ell(J)$ converges to $S \in \Gamma_\ell(J)$ iff $\|S_\alpha(a) - S(a)\| \rightarrow 0$. We call this the $\ell\kappa$ -topology. Obviously J is $\ell\kappa$ -dense in $\Gamma_\ell(J)$. Similarly one can define $r\kappa$ -topology on $\Gamma_r(J)$.

Lemma 3.1. If $J \subseteq A_0$ is a dense right ideal, then $b(\Gamma_\ell(J)) \subseteq M_\ell(A_0)$.

Proof This is to say that each bounded left multiplier of J extends uniquely to a left multiplier of A_0 , which follows from density of J in A_0 . $\square$

Definition 3.2. We say that $T \in L(A_0)$ is left-affiliated with A_0 if $z_T = T(I + T^*T)^{-\frac{1}{2}} \in M_\ell(A_0)$ and $z_T^* z_T < I$, or equivalently, if there is $z \in M_\ell(A_0)$ of norm one such that $\sqrt{1 - z_T^* z_T} A_0$ is dense in A_0 and $T(\sqrt{1 - z_T^* z_T} a) = za$. We denote the set of all these elements by $A_0^{\ell\eta}$. The elements right-affiliated with A_0 are defined similarly, and the set of all such elements is denoted by $A^{r\eta}$.

The following lemma is now trivial.

Lemma 3.2. For each dense right ideal $J \in A_0$, $\Gamma_\eta(J) \subseteq A_0^{\ell\eta}$. $\square$

Now let us come back to the special case of $A_0 = C^*(G)$ and $J = D_\ell^\infty$. Take any $X \in \mathcal{U}(\mathcal{G})$, then for each $a \in J$, $d_\ell u(X)a = \lim_{t \rightarrow 0} \frac{1}{t} (S_{\exp(tX)} - 1)(a)$, which means that $d_\ell u(X) = \ell\kappa\text{-lim } S_{\exp(tX)} - 1 \in \Gamma_\ell(J) \subseteq A_0^{\ell\eta}$. Therefore we have

Theorem 3.1. Let G be a Lie group and $\mathcal{G}$ be its Lie algebra. For each $X \in \mathcal{U}(\mathcal{G})$, $d_r u(X)$ and $d_\ell u(X)$ are respectively right-affiliated and left affiliated with $C^*(G)$. $\square$

We can get a better result using the generalized Stone's theorem, proved in previous section. Let A_0 be a C^* -algebra and G be a Lie group acting on A_0 through a strictly continuous representation $u : G \rightarrow M(A_0)$. Then

$$C^\infty(u) = \{x \in A_0 : g \mapsto u(g)x \text{ is } C^\infty\}$$

contains the dense subspace of A_0 spanned by all elements $u(f)x$, with $f \in C_{00}^\infty(G)$ and $x \in A_0$.

Proposition 3.1. With the above notation, to each $X \in \mathcal{G}$ there corresponds an element $\hat{X} \in \Gamma(A_{00})$ such that

$$e^{t\hat{X}} = u(e^{tX}) \quad (t \in \mathbb{R}).$$

Moreover $\hat{X}$ leaves $C^\infty(u)$ invariant and $[X, Y]^\wedge = [\hat{X}, \hat{Y}]$, when restricted to $C^\infty(u)$, for all $X, Y \in \mathcal{G}$.

Proof Take $X \in \mathcal{G}$ and put $U_t = u(e^{tX})$. Then Theorem 2.1 applies and gives $\hat{X} \in \Gamma(A_{00})$ with $U_t = e^{t\hat{X}}$ for each $t \in \mathbb{R}$ (here $\hat{X}$ is *skew* self adjoint). The rest is proved as in [HQV, 2.4]. $\square$

Corollary 3.1. *With the above notation, $\mathcal{U}(\mathcal{G}) \subseteq \Gamma(K(C^*(G)))$.*

Proof Just observe that G acts strictly continuously on $C^*(G)$ through the universal representation and apply above proposition. $\square$

Acknowledgement: This paper is part of the author's Ph.D. thesis in the University of Illinois at Urbana-Champaign under the supervision of Professor Zhong-Jin Ruan. I would like to thank him for his moral support and scientific guidance during my studies.

REFERENCES

- [Am] M. Amini; Locally compact pro- C^* -algebras , preprint.
- [HC] Harish-Chandra; Representations of a semisimple Lie group on a Banach space, I. Trans. Amer. Math. Soc. 75 (1953) 185–243.
- [HQV] J. Hollevoet, J. Quaegebeur, S. Van Keer; Stone's theorem in C^* -algebras , Quart. J. Math. Oxford Ser. (2) 43(170) (1992) 227-233.
- [LT] A. Lazar, D.C. Taylor: Multipliers of Pedersen's ideal . Mem. Amer. Math. Soc. 169 (1976).
- [Nel] E. Nelson; Analytic vectors, Ann. of Math. 70(2) (1959) 752-615.
- [Pd66] Gert K. Pedersen; Measure theory for C^* -algebras I – IV, Math. Scand. 19 (1966) 131-145; **22** (1968) 63-74; **25** (1969) 71-93; **25** (1969) 121-127.
- [Pd64] Gert K. Pedersen: Measure theory for C^* -algebras , Ph.D. Thesis.
- [Ph88a] N.C. Phillips: A new approach to the multipliers of Pedersen's ideal . Proc. Amer. Math. Soc. 104(3) (1988) , 861-867.
- [Ph88b] N.C. Phillips: Inverse limits of C^* -algebras and applications, Operator algebras and applications, Vol. 1, 127-185, London Math. Soc. Lecture Note Ser., 135 (1988) , Cambridge Univ. Press , Cambridge-New York.
- [Wr95] S.L. Woronowicz; C^* -algebras generated by unbounded elements, Rev. Math. Phys. 7(3) (1995) 481-521.
- [Wr91] S.L. Woronowicz; Unbounded elements affiliated with C^* -algebras and non-compact quantum groups , Com. Math. Phys. 136 (1991) 399-432.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, 106 WIGGINS ROAD, SASKATOON, SASKATCHEWAN, CANADA S7N 5E6, MAMINI@MATH.USASK.CA